

Unification of Gravity and Electro-Magnetism in Six Dimensions

Merab Gogberashvili

Andronikashvili Institute of Physics, 6 Tamarashvili Str., Tbilisi 380077, Georgia

E-mail: gogber@hotmail.com

February 1, 2008

Abstract

Recently by us was proposed the model where Einstein's equation on the brane was connected with Maxwell's multi-dimensional equations in pseudo-Euclidean space. Based on this idea unification of 4-dimensional gravity and electromagnetism in (2+4)-space is found. In this picture photon is massless in four dimensions and obtains large mass in extra (1+1)-space normal to the brane.

PACS numbers: 04.50.+h, 03.50.De, 98.80.Cq

One of the main ideas of the brane models is that Plank's scale M_{Pl} can be constructed with the fundamental scale M and the brane width ϵ [1]. In our previous paper [2] it was shown that in the brane approach Einstein's equations can be effective as well. Gravitation was connected with the geometry of the brane embedded in pseudo-Euclidean space, where solutions of 4-dimensional Einstein's equation could be constructed with the solutions of multidimensional Maxwell's equations. This approach is the inverse to the standard Kaluza-Klein picture, since the vector field is fundamental and gravity is obtained from the brane geometry. In this direction geometrical unification of different interactions is easier, since Dirac equation also can be derived from the constrained Yang-Mills Lagrangian [3]. The surface where Einstein's equations appear is not necessary to be 4-brane. For example, it can be 5-dimensional space with exponential warp factor needed to cancel cosmological constant [5]. Embedding of this 5-space into Euclidean (2+4)-space was found in [6].

The main idea of [2] was based on the fact that any n -dimensional Riemannian space can be embedded into N -dimensional pseudo-Euclidean space with $n \leq N \leq n(n+1)/2$ [4]. This means that the multidimensional 'tetrad' field h_α^A can be expressed as a derivative of embedded functions

$$h_\alpha^A = \partial_\alpha \phi^A . \quad (1)$$

Capital Latin letters $A, B, \dots$ numerate coordinates of embedded space, while Greek indices $\alpha, \beta, \dots$ numerate coordinates in four dimensions.

The metric of bulk flat space-time (where the brane with arbitrary geometry is embedded) can be transformed to the Gaussian normal coordinates

$$ds^2 = \eta_{ij} dx^i dx^j + g_{\alpha\beta}(x^i, x^\nu) dx^\alpha dx^\beta . \quad (2)$$

Here η_{ij} is the Minkowskian metric of the normal space to the brane and small Latin indices $i, j, \dots$ numerates extra $(N-4)$ coordinates. The induced metric on the brane is $g_{\alpha\beta}(0, x^\nu)$.

Embedding theory allows us to rewrite Hilbert's 4-dimensional action in terms of derivatives of the normal components ϕ^i of some multi-dimensional vector

$$S_g = -M_{Pl}^2 \int R \sqrt{-g} d^4 x = \eta_{ij} M_{Pl}^2 \int (\Box \phi^i \Box \phi^j - \partial_\alpha \partial_\beta \phi^i \partial^\alpha \partial^\beta \phi^j) \sqrt{-g} d^4 x , \quad (3)$$

where $\Box = \partial_\alpha \partial^\alpha$ is the 4-dimensional wave operator and g is the determinant in Gaussian coordinates.

In [2] it was shown that to the same form (3) can be transformed Maxwell's multi-dimensional action

$$S_A = -\frac{1}{4} \int F_{AB} F^{AB} d^N X = -\frac{1}{2} \int [\partial_A A_B (\partial^A A^B + \partial^B A^A) - 2 \partial_A A^A \partial_B A^B] d^N X , \quad (4)$$

where $F_{AB} = \partial_A A_B - \partial_B A_A$ and X^A are N-dimensional Cartesian coordinates. For this reduction we must transform X^A to the Gaussian coordinates (2) and insert in (4) the exact solution of Maxwell's equations for the brane

$$\begin{aligned} A^\alpha &= \left(\frac{2}{\pi\epsilon^2} \right)^{(N-4)/4} M \sum_i^{(N-4)} \exp \left[-\frac{(x^i)^2}{\epsilon^2} \right] \partial^\alpha \phi^i(x^\beta) , \\ A^i &= - \left(\frac{2}{\pi\epsilon^2} \right)^{(N-4)/4} \frac{2Mx^i}{\epsilon^2} \exp \left[-\frac{(x^i)^2}{\epsilon^2} \right] \phi^i(x^\beta) . \end{aligned} \quad (5)$$

Then we must integrate (4) by the extra coordinates x^i , remove boundary terms and as in [1] put

$$M_{Pl}^2 = \epsilon^{(N-4)} M^2 . \quad (6)$$

In the present paper we want to find 4-dimensional unification of gravity and Electro-Magnetism based on Maxwell's equations in 6-dimensional pseudo-Euclidean space with the signature $(2+4)$.

The $(2+4)$ -space is interest object for investigations. As it was found a long time ago Schwarzschild's metric can be embedded in $(2+4)$ -space [7]; also non-linear 15-parameter conformal group can be written as a linear Lorentz-type transformations there [8]. For compact Kaluza-Klein models $(2+4)$ -space was studied in [9] and in the context of brane models with non-factorizable geometry in [10].

In our previous paper [2] was considered only possible connection of Einstein's equations with multi-dimensional Maxwell's equation and was not received Maxwell's equations on the brane. In (5) there presents only derivatives of ϕ^i (which is connected with gravity) and it is clear that $F_{\alpha\beta}$ is zero. Now we want to insert in (5) also non-integrable part $a^\alpha(x^\beta)$ and identify it with the 4-dimensional vector-potential. So, in $(2+4)$ -space we want to rewrite (5) in the form

$$\begin{aligned} A^\alpha &= \sqrt{\frac{2}{\pi\epsilon^2}} \left[M \exp \left(-\frac{\tau^2}{\epsilon^2} \right) \partial^\alpha \phi^\tau(x^\beta) + M \exp \left(-\frac{\kappa^2}{\epsilon^2} \right) \partial^\alpha \phi^\kappa(x^\beta) + \exp \left(-\frac{\tau^2 + \kappa^2}{\epsilon^2} \right) a^\alpha \right] , \\ A^\tau &= -\sqrt{\frac{2}{\pi\epsilon^2}} \frac{2M\tau}{\epsilon^2} \exp \left(-\frac{\tau^2}{\epsilon^2} \right) \phi^\tau(x^\beta) , \\ A^\kappa &= -\sqrt{\frac{2}{\pi\epsilon^2}} \frac{2M\kappa}{\epsilon^2} \exp \left(-\frac{\kappa^2}{\epsilon^2} \right) \phi^\kappa(x^\beta) , \end{aligned} \quad (7)$$

where κ and τ are the normal to the brane space-like and time-like coordinates respectively.

Using the anzats (7) Maxwell's action (4) in $(2+4)$ -space takes the form

$$\begin{aligned} S_A &= -\frac{1}{4} \int F_{AB} F^{AB} d^6 X = \\ &= -M^2 \epsilon^2 \int R \sqrt{-g} d^4 x - \frac{1}{2\pi\epsilon^2} \int \exp \left[-\frac{2(\tau^2 + \kappa^2)}{\epsilon^2} \right] \left[f_{\alpha\beta} f^{\alpha\beta} + \frac{4}{\epsilon^4} (\tau^2 - \kappa^2) a_\alpha a^\alpha \right] \sqrt{-g} d^4 x d\tau d\kappa = \\ &= - \int \left(M_{Pl}^2 R + \frac{1}{4} f_{\alpha\beta} f^{\alpha\beta} \right) \sqrt{-g} d^4 x , \end{aligned} \quad (8)$$

where $f_{\alpha\beta} = \partial_\alpha a_\beta - \partial_\beta a_\alpha$. Latin indices are summing with the 6-dimensional Minkowskian metric tensor η_{AB} and Greek indices with the induced on the brane metric tensor $g_{\alpha\beta}$. So, 6-dimensional Maxwell's action on the brane is reduced to the 4-dimensional Einstein-Maxwell action. As in [10] the brane is placed at $\tau^2 = \kappa^2$ what means that it moves with the speed of light in transversal $(1+1)$ -space.

In Gaussian coordinates (2) Maxwell's 6-dimensional equations

$$\partial_A F^{AB} = 0 , \quad (9)$$

after inserting there the anzats (7) reduces to the system

$$\begin{aligned} \frac{1}{\sqrt{-g}} \partial_\alpha (\sqrt{-g} f^{\alpha\beta}) - \frac{4}{\epsilon^4} (\tau^2 - \kappa^2) a^\beta &= 0 , \\ \partial_\alpha a^\alpha &= 0 . \end{aligned} \quad (10)$$

One can see that the situation is similar to superconductors. On the brane ($\tau^2 = \kappa^2$) photon is massless and obtains the large mass $\sim 1/\epsilon$ in the direction normal to the brane.

At the end we want to note that even if we don't want to connect ϕ^i with gravity the ansatz (7) can be used for the trapping of photons on the brane in the (2+4)-space.

Acknowledgements: Author would like to acknowledge the hospitality extended during his visits at the Abdus Salam International Centre for Theoretical Physics where this work was done.

References

- [1] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett.* **B429**, 263 (1998);
I. Antoniadis, S. Dimopoulos and G. Dvali, *Nucl. Phys.* **B516**, 70 (1998).
- [2] M. Gogberashvili, hep-th/0203090.
- [3] Y. Takahashi, *J. Math. Phys.*, **24**, 1783 (1983);
V. A. Zhelnorovich, *Proc. Acad. Sci. USSR*, **311**, 590 (1990);
F. Reifler and R. Morris, *J. Math. Phys.*, **36**, 1741 (1995).
- [4] L. P. Eisenhart, *Riemannian Geometry* (New Jersey: Princeton University Press, 1949);
A. Friedman, *J. Math. Mech.*, **10**, 625 (1961); *Rev. Mod. Phys.*, **37**, 201 (1965).
- [5] M. Gogberashvili, hep-ph/9812296; *Mod. Phys. Lett.*, **A14**, 2024 (1999); *Europhys. Lett.*, **49**, 396 (2000); hep-ph/9908347;
L. Randall and R. Sundrum, *Phys. Rev. Lett.*, **83**, 3370 (1999); **83**, 4690 (1999).
- [6] M. Gogberashvili, gr-qc/0202061.
- [7] J. Rosen, *Rev. Mod. Phys.*, **37**, 204 (1965).
- [8] P. Dirac, *Ann. Math.*, **37**, 429 (1936); *Proc. Roy. Soc.*, **A155**, 447 (1936).
- [9] M. Pavsic, *Nuovo Cimento*, **B41**, 397 (1977);
R. L. Ingraham, *Nuovo Cimento*, **B46**, 1 (1978); **B46**, 16 (1978); **B46**, 217 (1978); **B46**, 261 (1978);
B47, 157 (1978); **B50**, 233 (1979); **B68**, 203 (1982); **B68**, 218 (1982).
- [10] M. Gogberashvili, *Phys. Lett.*, **B484**, 124 (2000);
M. Gogberashvili and P. Midodashvili, *Phys. Lett.*, **B515**, 447 (2001); hep-th/0111132.